

Physical Properties, Chemical Composition and Nutritional Quality Potentials of *Lonchocarpus sericeus* (Cube root) Seeds and Seed Oil

Running title

Nutritional quality potentials of *L. sericeus* seeds

Abstract

Physical properties, chemical composition and nutritional quality potentials of the seeds of *Lonchocarpus sericeus* (cube root) were investigated. The water absorption properties of the seeds were measured both gravimetrically as hydration capacity (0.02 ± 0.005 g/seed), and hydration index (0.06 ± 0.01) and volumetrically as swelling capacity (0.01 ± 0.00 mL/seed) and the swelling index ($1.2 \times 10^{-4} \pm 0.00$). A 100 g of seeds contained 17.53 ± 2.08 , 38.83 ± 4.16 , 3.18 ± 0.19 , 24.93 ± 3.21 , 29.96 ± 3.82 and 3.10 ± 0.89 g moisture, crude fibre, ash, crude fat, crude protein, and nitrogen free extract (NFE), respectively; 149.20 ± 0.24 kJ gross energy, 848 ± 16.1 , 722 ± 10.0 , 720 ± 9.1 , 230 ± 8.2 , and 224 ± 8.1 mg Fe, Mn, Zn, Cu and Na, respectively per 100 g seed sample. The seed also contained 3.14 ± 0.12 , 2.18 ± 0.32 and 4.30 ± 0.06 mg/g tannin, oxalate and saponin, respectively. The essential-to-total amino acid ratio (E/T, %) is 44.10%; average predicted protein efficiency ratio (PER) is 2.70, while the predicted biological value (BV) is 34.85. Elaidic acid is the most significant fatty acid (46.32%) found in the seeds with a total saturated and unsaturated fatty acids of 46.14% and 50.50%, respectively. The oil of *L. sericeus* is equally rich in bioactive compounds including tocopherols, sterols, phospholipids, tocotrienols and terpenoids with a concentration of 50.09, 479.13, 333.14, 1.34 and 3.33×10^{-4} mg/100g, respectively. *L. sericeus* seed would be a potential source of dietary energy, industrial oil, bioactive compounds and vegetable protein.

Keywords: Bioactive compounds, elaidic acid, vegetable oil, *Lonchocarpus sericeus*, micronutrients, underexploited legume.

Introduction

The problem of inadequate animal protein intake particularly in developing countries in the tropics due to the high cost and short supply is frequent [1, 2], and this is expected to escalate in the coming years due to unplanned population growth and insecurity fuelled by the insurgency. Nigeria currently spends over \$44 million annually on the importation of cowpea to bridge the gap between domestic production and cowpea consumption deficit of five hundred thousand tonnes.

Research into new seed crops for protein and oils for human and animals will conserve scarce foreign exchange, provide adequate nutrition that could alleviate the consequences of inadequate protein in diets, and also the various civilisation-related diseases including diabetes, cardiac diseases and obesity since many health-promoting roles have been ascribed to bioactive food components [3, 4]. Consequently, attention has been directed at sourcing alternative cheaper plant protein and seed oil to alleviate the problem of malnutrition in children and pregnant women, and also provide feeds for animals particularly in areas ravaged by drought and conflicts [5-8]. Several workers had acknowledged the suitability of *Cajanus cajan* (pigeon pea), *Vigna unguiculata* (cowpea) and *Glycine max* (soybean) as cheaper vegetable protein sources and alternatives to the expensive animal proteins [9, 10] while *Lannea kerstingii* [11], pomegranate, cherry, pumpkin [3, 12], chia [4], and garden cress (*Lepidium sativum* L.) seeds have also been documented for their excellent seed oils and bioactive components [13].

The present study reports the physical properties, chemical composition and nutritional quality potentials of *L. sericeus* (Cube root) seed. *L. sericeus* (Poir.) is a Leguminosae belonging to the subfamily *Papilionaceae*. It is commonly called cube root or Senegal lilac. Its seeds are usually oblong-kidney-shaped, reddish-brown, about 7 and 5 mm long and wide, respectively. It is native to several West African countries, including Nigeria, where its bark and roots have been explored for medicinal uses [14]. The plant grows wild, and its cultivation is supported by the climate and soil of the West African region [14, 15]. The seeds of *L. sericeus* had been reported to be abundant in flavonoids, particularly flavonol, quercetin, sugars, vitamins and bioactive principles including carotenoids and anthocyanins [16]; it is also

rich in insecticide rotenone [15]. The seed of *L. sericeus* remains unexploited, and if found nutritionally suitable, could serve as a partial replacement for cowpea as dietary protein and oil supply.

Materials and methods

Seed samples

In this study *L. sericeus* seeds were obtained from the biological garden of the Federal Polytechnic, Ilaro in South West Nigeria, (6.89°N, and 3.02°E) between March and April 2019. They were identified at the Forestry Research Institute of Nigeria (FRIN), Ibadan with herbarium number FHI 110133. A clean bulk, matured, healthy and unbroken seeds were used in the study. A portion of the bulk was removed for the determination of physical properties, while the remaining was reserved for chemical analysis. The latter portion was washed with deionized-distilled water, drained and dried in oven at 60°C; the seeds were pulverised in a Waring laboratory blender (MX-LBC15), sieved and kept in an airtight polythene bag for chemical analysis.

Physical properties of *L. sericeus* seeds

Seed weight, seed volume, seed density, hydration capacity, hydration index, swelling capacity and swelling index were determined as previously described [9, 17-19].

Briefly, the weight of 100 seeds was determined on an analytical balance and was recorded in grams (g). Seed volume was measured by transferring the previously weighed 100 seeds sample into a 100 mL measuring cylinder and 50 mL distilled water was added. Seed volume was expressed as an increase in volume per 100 seeds (mL/100seeds). Seed density was calculated by dividing the weight of 100 seeds sample by their volume measured with a 100 mL measuring cylinder. Hydration capacity was recorded as the gain in weight after an overnight soaking of 100 seed samples in distilled water; while hydration index was calculated as hydration capacity divided by the initial 100 seeds weight. Swelling capacity was the increase in the volume of 100 seeds sample after an overnight soaking in distilled water, and the swelling index was determined by dividing the swelling capacity with the initial seed volume and expressed as mL/seed. All the determinations were carried out in triplicates.

Proximate, mineral and anti-nutritional composition

Powdered seeds sample was used for the determination of moisture, Kjeldahl nitrogen, ether extract, and crude fibre and ash contents following the standard official methods [20]. Crude protein was estimated as %N multiplied by 6.25. Nitrogen free extractives (NFE) was determined by following equation (Eqn.) 1.

$$\%NFE = 100 - (\% \text{crude protein} + \% \text{crude fibre} + \% \text{crude fat} + \% \text{ash} + \% \text{moisture}). \text{ Eqn. 1}$$

Gross energy value of the seed was estimated in kilojoules by multiplying the crude protein, ether extract, and NFE by the factors of 16.7, 37.7 and 16.7, respectively [21]. Triplicate determinations were done for all the experiments.

Seed flour samples were also analysed for potassium, magnesium, calcium, sodium, manganese, iron, copper and zinc using Atomic Absorption Spectrophotometer (Buck 210 VGP) while phosphorus content was determined by colorimetric technique (Ammonium molybdovanadate) following standard official methods [20]. The oxalate, tannin, and saponin contents of seed flour were also determined by official methods [20].

Extraction of seed oil and determination of fatty acid composition

Soxhlet extraction of *L. sericeus* seed flour for oil was done with 400 mL n-hexane for 8 hrs. Solvent recovery was carried out by using a vacuum rotary evaporator (RE300 Bibby) at 40 – 50 °C. Determination of fatty acid composition was carried out using the official method [20] and modified method of Omar and Salimon [22]. One microliter of fatty acid methyl ester (FAME) was injected into a GC-FID (HP 6890) with an HP column (30 m x 0.25 mm x 0.25 µm particle size) with helium carrier gas. The injector temperature was 250 °C with split-less modes. The initial column temperature was set at 80 °C, and held for 1 min and ramped at 10 °C/min to 200 °C for 2 min then 20 °C/min to 300 °C for 8 min.

Six internal standards of saturated fatty acids (Palmitic, stearic, arachidic and lignoceric acids) and unsaturated fatty acids (Elaidic and oleic acids) were processed with n-hexane then their methyl esters were prepared, and analysed under the same conditions.

Bioactive components in the seed oil of *L. sericeus*

Tocopherols and tocotrienols in the seed oil was determined as described by Du and Ahn [23]. Sterols and terpenoids contents of the oils were evaluated using the modified methods of AOAC [24, 25]. The method of Raheja et al. [26] was used for analysis of phospholipids.

An HP 6890 GC-FID system was used for the analyses with split injection in the ratio of 20:1 and a detector temperature of 300°C. The inlet temperature was maintained at 290 °C with helium as the carrier gas for tocopherols and tocotrienols while nitrogen was the carrier gas for sterols, terpenoids and phospholipids. HPINNOWax column (30 m x 0.25 mm x 0.25 µm, particle size) was used for sterols and terpenoids analyses while HP-5ms UI with the same dimension was used for the studies of tocopherols, tocotrienols and phospholipids. The oven program had an initial temperature of 180°C for 0 min. The first ramp at 8°C/min to 260°C and second ramp at 2°C/min to 280°C was held constant for 13 minutes. The flow rate and hydrogen pressure were 2.5 mL/min and 22 psi, respectively, while data acquisition was achieved with HP ChemStation Rev. A 09.01 [1206] software.

Amino acid derivatization, analysis and protein quality evaluation

Determination of the amino acid content of *L. sericeus* was performed following modified AOAC and Danka et al. methods [20, 27]. A half gram of sample was weighed and added into 250 mL conical flask and defatted with 30 mL petroleum ether using Soxhlet extractor. The defatted sample was soaked in 30 mL of 1 M KOH solution and incubated for 48 h at 110°C in hermetically-closed borosilicate glass container. After the alkaline hydrolysis, pH of hydrolysate was adjusted to a pH range of 2.5 – 5.0. This was thereafter purified by cation-exchange solid-phase extraction. The amino acids in purified solution were derivatised with ethylchloroformate that was afterwards removed by passing streams of nitrogen gas. Aliquots of amino acids derivatives dissolved in dichloromethane were analysed by gas chromatography equipped with pulsed flame photometric detector (GC-PFPD). One microliter of concentrate was injected into the gas chromatograph (HP 6890) with HP-5MS column (30 m x 0.25 mm, 0.25 µm particle size) for individual amino acid peaks. The initial temperature of the hydrogen carrier gas and column was 60 °C. It was ramped at 8°C for 20 min and held constant for 2 min and then increased to 12°C/min for 6 min.

Amino acids standard solutions were repeatedly analysed for five times and the calibration curves obtained had correlation coefficients ≥ 0.9992 .

Subsequently, the quality of seed protein was evaluated by:

- a. Calculating the proportion of essential amino acids (E) to the total amino acids

$$(T) \times 100, \left(\frac{E}{T} \times 100\right) \quad \text{Eqn. (2)}$$

- b. Computing essential amino acids index (EAAI) using Eqn. (3)

$$\text{EAAI} = 10^{\log \text{EAA}} \quad \text{Eqn. (3)}$$

$$\text{where } \log \text{EAA} = 0.1[\log \left(\frac{a_1}{a_s} \times 100\right) + \log \left(\frac{a_2}{a_s} \times 100\right) \dots + \log \left(\frac{a_n}{a_{ns}} \times 100\right)]$$

$$a_1 \dots \dots a_n = \text{Amino acids content of the sample}$$

$$a_s \dots \dots a_{ns} = \text{Amino acids values from the reference i.e. whole egg}$$

Herein, amino acids values from the reference are: Ile = 4.00; Leu = 7.04; Lys = 5.44; Met + Cys = 3.52; Trp = 0.56; Val = 4.76; Phe + Tyr = 6.08; Thr = 4.00; and His = 2.40.

- c. Biological Value (BV), was estimated from EAAI

$$\text{BV} = 1.09(\text{EAAI}) - 11.73 \quad \text{Eqn. (4)}$$

- d. The protein efficiency ratio (PER) was predicted from the amino acid content according to the equation

$$\text{PER} = -0.468 + 0.454(\text{Leu}) - 0.105(\text{Tyr}) \quad [28] \quad \text{Eqn. (5)}$$

Results and discussion

Physical properties

The physical properties of *L. sericeus* seeds are presented in Table 1. The weight of 100 seeds of *L. sericeus* is similar to that of *Lupinus albinus* (28.65±0.28 g [18]) but slightly heavier than *Lathyrus maritimus* seeds (25.27±0.05 g) [17]. It is about three times heavier than the seeds of *C. cajan* (10.57±1.029 g) [9]; but lighter than the seeds of *Parkia roxburghii* (53.00±4.00 g [19]). The weight of *L. sericeus* as determined in this study could be classified alongside the medium market class of 25-40 g/100 seeds, where most common bean (*Phaseolus vulgaris* L.) species are grouped. The market classes are grouped as small ≤ 25 , medium

25-40, and large ≥ 40 g/100 seeds [29]. The seed volume (mL/100 seeds) of 24.75 ± 0.96 followed a similar pattern with the seeds earlier stated. This is reflected in the density of the seed as it is similar to the density of all the seeds mentioned previously except that of *P. roxburghii* (0.45 ± 0.05).

The hydration capacity (0.02 ± 0.00 g/100 seeds), hydration index (0.06 ± 0.008), swelling capacity (0.011 ± 0.00 mL/seed) and swelling index ($1.2 \times 10^{-4} \pm 0.00$) as presented in Table 1 showed comparable values with those of matured *L. maritimus* [7] and *C. cajan* [9]. Arising from their low physical characteristic values, *L. sericeus* seeds would have poor swelling properties that could make them unsuitable for some domestic applications that include the preparation of puddings, soups and sauces [29]. These characteristics may not be unconnected with the fact that *L. sericeus* seeds have a glossy and hard coat, therefore, making them be impermeable and not easily hydrated. Consequently, it may take a longer time for the seeds to germinate or cook as cooking time positively correlates to seed physical properties such as density, swelling index, hydration capacity and swelling capacity and seed coat characteristic; these qualities influence the inclination of consumers and processors of seed legumes [17, 30].

Table 1. Physical Properties of the seed of *L. sericeus*

Parameter	Mean \pm SD
Seeds weight (g/100 seeds)	28.08 ± 0.45
Seed volume (mL/100 seeds)	24.75 ± 0.96
Seed density (g/mL)	1.14 ± 0.04
Hydration capacity (g/100seeds)	0.02 ± 0.00
Hydration index	0.06 ± 0.01
Swelling capacity (mL/seed)	0.011 ± 0.00
Swelling index	$1.2 \times 10^{-4} \pm 0.00$

Proximate, mineral and antinutrient composition of *L. sericeus*

The chemical composition of *L. sericeus*, as shown in Fig.1 compares well with those of cowpea (*V. unguiculata*) and scarlet runner bean (*Phaseolus coccineus*) [31]. The composition of different varieties of cowpea and scarlet runner bean previously reported in g/100 g contained moisture (0.4-1.8), ash (3.61-4.11), crude fat (2.13 – 7.53), and crude protein (7.53-52.61) while the results from our present study

showed *L. sericeus* to be higher in crude protein than cowpea (7.53 ± 0.02) but less than scarlet runner bean (51.14 ± 0.0); however, crude protein value from the present work is greater than the 28.03% reported for *L. sericeus* in a previous study [32]. In addition, results from this study also showed higher values than crude fat ($3.45 \pm 0.06\%$), crude fiber ($6.85 \pm 0.07\%$), crude protein ($23.45 \pm 0.62\%$), ash ($3.55 \pm 0.04\%$), but lower in carbohydrate ($58.89 \pm 0.44\%$) reported for cowpea cultivars from Sri Lanka [33] and still lesser than the average of 62.32 ± 2.18 g/100 g for different accessions of *Mucuna* beans [34]. The crude protein and ash contents of *L. sericeus* compare with 29.83 ± 1.15 and 3.44 ± 0.3 g/100 g, respectively, reported by Mohan and Kala, 2010 [35] for *M. pruriens* (Velvet bean) but richer in lipid and crude fibre contents. *L. sericeus* has higher protein, fibre and fat content than black turtle bean (*P. vulgaris*) that had 23.7 ± 0.15 protein, 3.3 ± 0.01 crude fibre and 15.2 ± 0.05 fat [36]. The crude ether extract of 24.93 ± 3.21 g/100 g for *L. sericeus* is very high compared to the 22.5 and 1.3 g/100 g documented by El-Shemy et al. [37] for soybean and faba beans, respectively; it is, however, about half the 45.41-48.14 g/100 g reported by many workers for *Arachis hypogea* [38, 39]. *L. sericeus* can invariably be classified as an oil-bearing seed since it contains oil in the range of those classified as oil-rich seeds. Comparing the results obtained in this study with soya beans seeds, *L. sericeus* had higher crude fibre but lower crude protein, ash and nitrogen-free extract than soya bean seeds [40].

The gross energy value of 149.2 ± 0.2 kJ/100 g (Table 2) obtained in the present study is not unexpected because of its high crude ether extract. The value compares well with 168.4 ± 0.2 kJ/100 g reported for different accessions of *Mucuna* beans [34] but less than 1597.6 - 1654.0 and 1600.6 kJ /100 g for *V. aconitifolia* and *V. unguiculata*, and *C. catharica*, respectively [41, 42].




Fig 1. Proximate composition (g/100g) of the seeds of *L. sericeus*

The results of mineral composition of *L. sericeus* in mg/100 g is as shown in Fig.2. *Lonchocarpus sericeus* is very rich in Na (224.8±8.10 mg/100 g) but poor on other macroelements when compared to cowpea that had 1062, 745.4 and 416.4 mg/100 g Ca, K and Mg, respectively [6]; and 443.0±12.17, 1417.5±21.65, 101.7±7.20 and 109.8±3.81 for P, K, Ca and Mg, in that order for *M. pruriens* [35]. Equally abundant in *L. sericeus* seeds are micronutrients including copper (230.0±8.21), zinc (720.0±9.06), manganese (722.0±10.04) and iron (848.0±16.09 mg/100 g). The amount of sodium in *L. sericeus* and the other micronutrients as mentioned earlier far exceeded what had been reported for most legumes. The seed also contained more iron and zinc than newly developed chickpea (*Cicer arietinum* L) varieties [43], and more sodium than in the chickpea reported by Simsek et al. [44] thus making *L. sericeus* seeds an excellent source for sodium and the micronutrients it contained. The Ca/P, and Na/K ratios obtained in this study (Table 2) may promote loss of calcium and high blood pressure [31]; and also, since the excess of one mineral has

been reported to decrease the absorptivity and proper utilisation of another for good nutrition [45], potassium supplementation may be required to correct the Na:K imbalance in the seeds of *L. sericeus* while the Ca:P revealed high value of phosphorus, therefore there may be a need to complement *L. sericeus* with seed grains rich in calcium.




Fig 2. Mineral composition (mg/100 g) of the seed of *L. sericeus*

The results of tannins, oxalates, and saponin contents of *L. sericeus* are as shown in Table 2. They are 3.14 ± 0.12 , 2.18 ± 0.32 , and 4.30 ± 0.06 in mg/g, respectively. These values are higher compare to those reported by Soetan et al. [46] for the seeds of *P. biglobosa* (tannin, 0.51 ± 0.00 ; oxalate, 0.47 ± 0.01 ; and saponin, 0.92 ± 0.00 mg/g) and approximately two times the values of the same parameters reported by Carew et al. [47] for *M. puriens* seeds. However, the values presented in this study are generally lower than those reported for most commonly consumed legumes including red kidney beans, lentils, white beans, Anasazi beans, and blackeyed pea [48]. From the concentrations of saponin, tannin, and oxalate obtained in this study, it is not likely they will affect the nutritional potentials of *L. sericeus* to any significant extent and neither will they pose any undesirable effects on its utilization.

Table 2. Gross energy, minerals ratio and anti-nutritional composition of the seed of *L. sericeus*

Gross energy kJ/100 g	149.20±0.24
Na/K	23.86
Ca/P	0.26
Anti-nutritional constituents (mg/g)	
Tannins	3.14±0.12
Oxalate	2.18±0.32
Saponin	4.30±0.06

Fatty acid composition of *L. sericeus*

The fatty acid composition of *L. sericeus* seed is indicated in Table 3 with more unsaturated fatty acids than saturated. The seed is very rich in dietary saturated fatty acids with palmitic acid (22.88 g/100 g) been the most abundant. Elaidic acid (46.32 g/100 g), a trans - ω^9 fatty acid is the most abundant in *L. sericeus*; though not an essential fatty acid like ω^3 and ω^6 fatty acids, ω^9 fatty acids protect against the possibility of metabolic syndrome and cardiovascular disease risk factors. They have been shown to increase high-density lipoprotein cholesterol and decrease low-density lipoprotein cholesterol; they also help reduce plaque buildup in the arteries that could pre-dispose an individual to a heart attack or a stroke [49]. ω^9 fatty acids provide energy, besides they are structural components of cell membranes, and contribute desirable texture and tastiness to foods [50]. The ratio of unsaturated to saturated fatty acids (1.1) obtained in this study reveals that consumption of *L. sericeus* oil will supply healthy fatty acids and can also overcome the negative effects associated with fatty acids [19].

Table 3. Fatty acid composition* of *L. sericeus* seed oil in g /100 g lipid

Saturated fatty acids	
Palmitic acid	22.88
Stearic acid	9.04
Behenic acid	7.84
Arachidic acid	2.65
Lignoceric acid	2.50
Margaric acid	1.23
Unsaturated fatty acids	
Elaidic acid (18:1)	46.32
Oleic acid	4.03
Cinnamic acid	0.15
Others	
Warfarin	0.78
Megastigmatrienone	0.56
Cuparene	0.20
Alpha-phenyl-o-xylene	1.81
Total saturated	46.14
Total unsaturated	50.50
Unsaturated / Saturated	1.10

Bioactive components in the seed oil of *L. sericeus*

The seed oil of *L. sericeus* contained all the eight different isomers of vitamin *E* with γ -tocopherol been the most abundant at about 89% similar to that of soya bean oil (88%) and higher than 10.5, 58, and 46% for Brazil nut (*Bertholletia excelsa*), olive and walnut oils, respectively [51, 52]. The total concentration of α - and γ -tocopherols in the oil was 50.09 mg/100 g, while the total tocol composition was 51.43 mg/100 g as shown in Table 4. The value is higher than the 42.54 mg/100 g total tocopherol reported by Nehdi [51] for cold pressed olive oil; while it compared with the 55 mg/100 g total tocopherol in coconut oil; and, however, higher than 33 mg/100 g for palm kernel oil [53]. Tocotrienols, though more potent in antioxidant activity [54] are reportedly less abundant in plant products compared to tocopherols [55], and this is clearly demonstrated in the present study. Due to its high levels of γ -tocopherol, the oil can find use in cosmetic and pharmaceutical formulations by adding value and stability to those products because of their inherent antioxidant capacity; and in some instances improve their emollient effects.

Taraxerol, α -amyrin, β -amyrin, luperol and bauerenol acetate were confirmed in the seed oil in trace amounts of 0.333 mg/100 g as shown in Table 4. α - and β -amyrin isolated from *Alstonia boonei* stem bark reportedly had profound anti-inflammatory activities in rodents [56]. Ebajo et al. [57] also isolated these terpenoids from *Hoya multiflora* (shooting-star hoyo) stem and leaves but did not report any biological activity on the isolates. De Amorim and co-workers, however, isolated the same principle from the leaves of *Pseudobrickellia brasiliensis* used as painkillers and in the treatment of inflammation [58]. The presence of terpenoids in the oil sample suggests that it could be useful as an analgesic and anti-inflammatory agent. *Lonchocarpus sericeus* seed oil is rich in phosphatidylcholine (230.37 mg/100 g) accounting for 69.15% of total phospholipids. Phosphatidylinositol and phosphatidylethanolamine were the next most abundant as indicated in Table 4. The total phospholipids in *L. sericeus* was 333.14 mg/100 g. Phospholipids have been established to improve the antioxidant properties of oils because of the synergism between them and tocopherols [59]. They also through their phosphate groups, chelate pro-oxidant metals and act as an oxygen barrier between oil and air interfaces [60]. Hidalgo et al. [61] used the Rancimat method to demonstrate that the addition of mixtures of amino acids to phospholipids further enhanced their antioxidant properties. Phosphatidylcholine and phosphatidylserine have respectively been reported in the treatment of liver conditions and the prevention of brain deterioration [62].

Sterols in the present study were found in amounts ranging from 302-310, 105-107, 52-78 and 12-16 mg/100 g in that order for sitosterol, campesterol, stigmasterol and Δ^5 -avenasterol, respectively with sitosterol been the most prevalent and accounted for over 60%. Other steroids determined in trace quantities include cholesterol, cholestanol and ergosterol. The total sterol contents for *L. sericeus* was 479.13 mg/100 g. The campesterol content exceeded the 4% maximum acceptable limit against the total sterol content for olive oils with stigmasterol also exceeding limits [63]. The sterol composition of the seed oil is higher than the 146.75 - 247.68 mg/100 g reported for different Tunisian olive oils shown to have good nutritional characteristics [64], and also the 204.0-248.0 mg/100 g for acorn fruit oils [65]. Sterols from plants, like other sterols are critical constituents of cellular membranes where they regulate fluidity and permeability [66, 67]. They have been reported to have the ability to lower blood serum cholesterol in

hypercholesterolemic and diabetic patients and healthy human volunteers [68]; they are also as effective as many cholesterol-lowering drugs [69]. The consumption of vegetable sterols has been associated with a lower risk of myocardial infarction, lowering LDL cholesterol even when combined with non-fat matrices. Additionally, they may moderate biomarkers of oxidative stress and inflammation and modulate the development of atherosclerosis.

Table 4. Bioactive components in the seed oil of *L. sericeus*

Bioactive compounds (mg/100 g)		<i>L. sericeus</i>
Tocopherols	α -tocopherol	3.83
	γ -tocopherol	45.94
	β -tocopherol	5.98×10^{-5}
	δ -tocopherol	3.19×10^{-1}
		50.09
Tocotrienols	α -tocotrienols	4.33×10^{-1}
	γ -tocotrienol	8.34×10^{-1}
	β -tocotrienol	7.36×10^{-2}
	δ -tocotrienol	1.48×10^{-4}
		1.34
Terpenoids	Taraxerol	3.18×10^{-4}
	α -amyrin	1.35×10^{-2}
	β -amyrin	3.09×10^{-1}
	Luperol	1.74×10^{-3}
	Baurenol acetate	8.59×10^{-3}
		3.33×10^{-1}
Phospholipids	Phosphatidylethanolamine	26.31
	Phosphatidylcholine	230.37
	Phosphatidylserine	4.59
	Lysophosphatidylcholine	2.29
	Phosphatidylinositol	69.58
		333.14
Sterols	Cholesterol	5.05×10^{-6}
	Cholestanol	4.67×10^{-4}
	Ergosterol	1.57×10^{-5}
	Campesterol	107.48
	Stig-masterol	52.30
	Δ^5 -avenasterol	16.80
	Sitosterol	302.55
		479.13

Amino acid composition of *L. sericeus*

The amino acid profile of *L. sericeus* in g/100 g of protein is presented in Table 5. The concentrations of glutamic acid (18.22), leucine (7.68), aspartic acid (7.14), proline (6.10), arginine (5.99), valine (5.47), and lysine (5.35) were high while the concentrations of methionine and cystine were the lowest (2.13 and 1.84 g/100 g), however, tryptophan was not detected; while leucine and arginine at 7.68 and 5.99 g/100 g, respectively were the most abundant essential amino acids. The ratio of essential amino acids to total obtained for *L. sericeus* was 44.10% and is above the 36% considered adequate for an ideal protein [70]. The protein efficiency ratio (PER) of 2.70 obtained in this study is higher than 1.87, 1.21, and 1.82 reported for *P. biglobosa*, *V. unguiculanta*, and *C. cajan*, respectively [71, 72]. The essential amino acid index (EAAI) of the seeds was 42.73, and the biological value (BV) of 34.85 was computed from it. The BV of 46.68 and 48.31 reported for roasted and pressure-cooked seeds of *Canavalia cathartica*, [7] and 63.35 for Bambara bean (*V. subterranea*) [73] are greater than 34.85 obtained in the present study; however this value is more than the 22.56 for *P. biglobosa* flour [71]. The contents of essential amino acids of *L. sericeus* especially leucine, valine, threonine and histidine were higher than those of the FAO/WHO that were found to be sufficient for an ideal protein while lysine is at a comparable level with recommended pattern [74]. The values of essential amino acids such as Leu, Val, Arg and Lys reported in this study are higher when compared to those found in soybean seeds [75].

Table 5. Amino acid content in g/100 g seed of *L. sericeus*

Essential amino acids	
Leucine	7.68
Valine	5.47
Isoleucine	3.73
Threonine	4.49
Histidine	3.02
Lysine	5.35
Methionine	2.13
Cystine	1.84
Phenylalanine	4.00
Arginine	5.99
Non-essential amino acids	
Glycine	4.88
Alanine	4.28
Serine	4.28
Proline	6.10
Aspartate	7.14
Glutamate	18.22
Tyrosine	3.09
Total	91.66
Protein quality evaluation	
E/T, %	44.49
PER	2.70
BV	34.85
EAAI	42.73

Conclusions

This study has reported chemical composition, physical properties and nutritional potentials of *L. sericeus*. The seeds were shown to be high in crude protein and fat contents but low in carbohydrate content. The amounts and quality of its oil and protein compared well to those for soybean and other common staples, it therefore clearly shows promise as a protein supplement for low-protein foods and feeds such as cereal grains. As oil-rich seed with a good fatty acid profile, its oil can be exploited for human and animal nutrition after further investigation into its characteristics and the oil could be exploited for industrial purposes. The seed has an interesting composition in macro and micronutrients, and also minimal in antinutrients thus having the potential to cover human nutritional requirements. With increasing population growth and continuous depletion of available food resources, *L. sericeus* represents a potential food source, which could reduce protein shortages and expand available food options and varieties, generate income, and afford

farmers the opportunity to spread risk in times of crop failure. However, further studies into its possible toxic and antinutrient factors such as L-Dopa and trypsin inhibitors, and digestibility is still required.

Acknowledgement

The authors are grateful to the anonymous reviewers for their useful comments

References

- [1] HC Schonfeldt, and NG Hall. Dietary protein quality and malnutrition in Africa. *Br. J. Nutr.* 2012; **108**, S69–S76.
- [2] S Ghosh. Assessment of protein adequacy in developing countries: Quality matters. *Food Nutr. Bull. Suppl.* 2013; **34(2)**, 244-246.
- [3] F Siano, MC Straccia, M Paolucci, G Fasulo, F Boscaino and MG Volpe. Physico-chemical properties and fatty acid composition of pomegranate, cherry and pumpkin seed oils. *J. Sci. Food Agric.* 2016; **96**; 1730–1735
- [4] B Kulczynski, J Kobus-Cisowska, M Taczanowski, D Kmiecik and A Gramza-Michałowska. The chemical composition and nutritional value of Chia seeds—Current state of knowledge. *Nutrients.* 2019, **11**, 1242. DOI: 10.3390/nu11061242
- [5] D Montesano, F Blasi, MS Simonetti, A Santini and L, Cossignani. Chemical and nutritional characterization of seed oil from *Cucurbita maxima* L. (var. Berrettina) Pumpkin. *Foods.* 2018, **7**, 30. DOI: 10.3390/foods730030
- [6] OC Adebooye and S Singh. Effect of cooking on the profile of phenolics, tannins, phytate, amino acid, fatty acid and mineral nutrients of whole-grain and decorticated vegetable cowpea (*Vigna unguiculata* L. Walp). *J. Food Qual.* 2007; **30**, 1101-1120.
- [7] KR Sridhar, S Seena, AB Arun and C-C Young. Effect of roasting and pressure-cooking on nutritional and protein quality of seeds of mangrove legume *Canavalia cathartica* from southwest coast of India. *J. Food Compos. Anal.* 2006; **19**, 284–293.
- [8] K Becker, P Siddhuraju and PSH Makkar. Chemical composition, protein fractionation, essential amino acid potential and antimetabolic constituents of an unconventional legume, Gila bean (*Entada phaseoloides* Merrill) seed kernel. *J. Sci. Food Agric.* 2001; **82(2)**, 192–202.
- [9] RA Oloyo and SS Akoja. Studies on the intrinsic physico-chemical properties of pigeon pea (*Cajanus cajan*) seed flour. *Trop. Agric. Res. Ext.* 2005; **8**, 78-83.
- [10] SPC Acuña, JHG González and IDA Torres. Physicochemical characteristics and functional properties of vitabosa (*Mucuna deeringiana*) and soybean (*Glycine max*). *Ciênc Tecnol Aliment.* 2012; **32(1)**, 98-105.
- [11] JT Ouilly, P Bazongo, A Bougma, N Kabore, AM Lykke, A Ouedraogo, and IHN Bassole. Chemical composition, physicochemical characteristics, and nutritional value of *Lannea kerstingii* seeds and seed oil. *J. Anal. Methods Chem.* 2017, DOI: 10.1155/2017/2840718
- [12] M Raczyk, A Siger, E Radziejewska-Kubzdela, K Ratusz and M Rudzińska. Roasting pumpkin seeds and changes in the composition and oxidative stability of cold-pressed oils. *Acta Sci. Pol. Technol. Alim.* 2017; **16(3)**, 293–301.
- [13] CS Singh and VK Paswan. The potential of Garden Cress (*Lepidium sativum* L.) seeds for development of functional foods. *Adv. Seed Biol.* 279-294. 2017, DOI: 10.5772/intechopen.70355.
- [14] FAO Food and Agriculture Organization of the United Nations, (1987). Some medicinal forest plants of Africa and Latin America, **FAO forestry paper No. 67**, Food and Agriculture Organization of the United Nations Viale delle Terme di Caracalla, Rome.
- [15] MM Iwu. Pharmacognostical profile of selected medicinal plants from: Handbook of African Medicinal Plants CRC Press, Florida 2014, p. 251-252.
- [16] OA Oyedeji, L Azeez, SO Adewuyi, BG Osinfade and MO Bamimore. Flavonoid profile, anthocyanin, carotenoid, sugar and vitamin compositions of *Lonchocarpus sericeus* seeds. *Afr. J. Food Sci. Technol.* 2015, **6(5)**, 131-135.

- [17] F Shahidi, UD Chavan, AK Bal and DB McKenzie. Physico-chemical properties and nutrient composition of beach pea (*Lathyrus maritimus* L.). *Food Chem.* 1999; **66**, 43-50.
- [18] SA Emire and H Tizazu. Chemical composition, physicochemical and functional properties of Lupin (*Lupinus albus*) seeds grown in Ethiopia. *Afr. J. Food, Agric., Nutr. Dev.* 2010; **10(8)**, 3029-3045
- [19] P Siddhuraju and A Sathya. Effect of processing methods on compositional evaluation of underutilized legume, *Parkia roxburghii* G. Don (yongchak) seeds. *J. Food Sci. Technol.* 2015; **52(10)**, 6157–6169
- [20] AOAC Official Methods of Analysis, 1990. 15th edn., The Association of Official Analytical Chemists, Arlington, USA.
- [21] S Ekanayake, ER Jansz and BM Nair. Proximate composition, mineral and amino acid content of mature *Canavalia gladiata* seeds. *Food Chem.* 1999; **66(1)**, 115-119.
- [22] TA Omar and J Salimon. Validation and application of a gas chromatographic method for determining fatty acids and trans fats in some bakery products. *J. Taibah Uni. Sci.* 2013; **7**, 56–63
- [23] M Du and D U Ahn. Simultaneous analysis of tocopherols, cholesterol and phytosterols using gas chromatography. *J. Food Sci.* 2002; **67(5)**, 1696-1700.
- [24] AOAC International. 1996. Official methods of analysis of AOAC international. 16th ed. Gaithersburg, Md.: AOAC Intl.
- [25] AOAC International. 1996. Official methods of analysis of AOAC international. 16th ed. Gaithersburg, Md.: AOAC Intl.
- [26] RK Raheja, C Kaur, A Singh and IS Bhatia. New colorimetric method for the quantitative estimation of phospholipids without acid digestion. *J. Lipid Res.* 1973; **14**, 695-697.
- [27] PO Danka, DT Dobrina and VI Kalin. Simultaneous identification and determination of total content of amino acids in food supplements – tablets by gas chromatography. *Asian J. Pharm. Clin. Res.* 2012; **5(2)**, 57-68
- [28] F Shahidi, UD Chavan and DB McKenzie. Functional properties of protein isolates from beach pea (*Lathyrus maritimus* L.). *Food Chem.* 2017; **4**, 177-187.
- [29] N Nciri, F El-Mhamdi, H Ismail, A Mansour and F Fennira. Physical properties of three white bean varieties (*Phaseolus vulgaris* L.) grown in Tunisia. *J. Appl. Sci. Agric.* 2014; **9(11)**, 195-200.
- [30] PZ Bassinello, B-S Siqueira, RP Vianello and KF Fernandes. Hardness of carioca beans (*Phaseolus vulgaris* L.) as affected by cooking methods. *LWT -- Food Sci Technol.* 2013; **54**, 13-17.
- [31] MO Aremu, O Olaofe and TE Akintayo. Compositional evaluation of cowpea (*Vigna unguiculata*) and scarlet runner bean (*Phaseolus coccineus*) varieties grown in Nigeria. *J. Food, Agric. Environ.* 2006; **4(2)**, 39 - 43
- [32] IE Ezeagu, CC Metges, J Proll, KJ Petzke and AO Akinsoyinu. Chemical composition and nutritive value of some wild-gathered tropical plant seeds. *Food Nutr. Bull. Suppl.* 1996; **17(3)**, 275-278.
- [33] R Liyanage, OS Perera, P Wethasinghe, BC Jayawardana, JK Vidanaarachchi and R Sivakanesan. Nutritional properties and antioxidant content of commonly consumed cowpea cultivars in Sri Lanka. *J. Food Leg.* 2014; **27(3)**, 215-217
- [34] IE Ezeagu, B Maziya-Dixon and G Tarawali. Seed characteristics and nutrient and antinutrient composition of 12 *Mucuna* accessions from Nigeria. *Trop. Subtrop Agroecosyst.* 2003; **1**, 129 – 139.
- [35] VR Mohan and BK Kala. Chemical composition and nutritional evaluation of lesser known pulses of the genus, *Mucuna*. *Adv. Bio Res.* 2010; **1(2)**, 105-116.
- [36] MO Aremu SS Audu and L Lajide. Influence of traditional processing methods on the nutritional composition of black turtle bean (*Phaseolus vulgaris* L.) grown in Nigeria. *Int. Food Res. J.* 2013; **20(6)**, 3211-3220.
- [37] H El-Shemy, E Abdel-Rahim, O Shaban, A Ragab, E Carnovale and K Fujita. Comparison of nutritional and antinutritional factors in soybean and fababean seeds with or without cortex. *J. Soil Sci. Plant Nutr.* 2000; **46(2)**, 515-524
- [38] OA Shokunbi, ET Fayomi, OS Sonuga and GO Tayo. Nutrient composition of five varieties of commonly consumed Nigerian groundnut (*Arachis hypogaea* L.). *Grasas Aceites.* 2012; **63**, 90-96
- [39] G Eshun, EA Amankwah and J Barimah. Nutrients content and lipid characterization of seed pastes of four selected peanut (*Arachis hypogaea*) varieties from Ghana. *Afr. J. Food Sci.* 2013; **7(10)**, 375-381
- [40] JE Van Eys, A Offner and A Bach. Chemical Analysis. Manual of Quality Analysis for Soybean Products in the Feed Industry. *Proc. Annu. Meet. Am. Soybean Assoc.* American Soybean Association. (2004)

- [41] VR Mohan and PT Soris. Chemical analysis and nutritional assessment of two less known pulses of genus *Vigna*. *Trop. Subtrop Agroecosyst.* 2011; **14**,473 – 484.
- [42] KR Sridhar, B Bhagya, S Seena, C-C Young, AB Arun and KV Nagaraja. Nutritional qualities and in vitro starch digestibility of ripened *Canavalia cathartica* beans of coastal sand dunes of southern India. *EJEAFChe.* 2006; **5(2)**, 1241-1252.
- [43] S Sharma, N Yadav, A Singh and R Kumar. Nutritional and antinutritional profile of newly developed chickpea (*Cicer arietinum* L) varieties. *Int. Food Res. J.* 2013; **20(2)**, 805-810
- [44] S Simsek, EN Herken and M Ovando-Martinez. Chemical composition, nutritional value and *in vitro* starch digestibility of roasted chickpeas. *J. Sci. Food Agric.* 2015; **96(8)**, 2896 –2905.
- [45] A Iqbal, IA Khalil, N Ateeq and MS Khan. Nutritional quality of important food legumes. *Food Chem.* 2006; **97**,331–335.
- [46] KO Soetan, AS Akinrinde and SB Adisa. Comparative studies on the proximate composition, mineral and anti-nutritional factors in the seeds and leaves of African locust bean (*Parkia biglobosa*). *Anna. Food Sci. Technol.* 2014; **15(1)**, 70-74.
- [47] LB Carew and AG Gernat. Use of velvet beans, *Mucuna* spp., as a feed ingredient for poultry: a review. *World's Poultry. Sci. J.* 2006; **62(01–14)**, 1314
- [48] BD Oomah, G Loarca-Pina and R Campos-Vega. Minor components of pulses and their potential impact on human health. *Food Res. Int.* 2010; **43**, 461–482.
- [49] LG Gillingham, S Harris-Janz and PJ Jones. Dietary monounsaturated fatty acids are protective against metabolic syndrome and cardiovascular disease risk factors. *Lipids.* 2011; **46(3)**, 209-228.
- [50] JE Hunter, J Zhang and PM Kris-Etherton. Cardiovascular disease risk of dietary stearic acid compared with trans, other saturated, and unsaturated fatty acids: a systematic review. *Am. J. Clin. Nutr.* 2010; **91**, 46 – 63
- [51] IA Nehdi, H Sbihi, CP Tan, H Zarrouk, MI Khalil and SI Al-Resayes. Characteristics, composition and thermal stability of Acacia Senegal (L.) Wild. *Ind. Crops Prod.* 2012; **36**, 54-58.
- [52] T Chunhieng, A Haidi, D Pioch, J Brohier and D Montet. Detailed study of Brazil nuts (*Bertholletia excelsa*) oil micro-compounds: phospholipids, tocopherols and sterols. *J. Braz. Chem. Soc.* 2008; **19(7)**, 1374–1380.
- [53] K Ellwood, DA Balentine, JT Dwyer, JW Erdman Jr., PC Gaine and CL Kwik-Urbe. Considerations on an approach for establishing a framework for bioactive food components. *Adv Nutr.* 2014; **5**, 693–701.
- [54] EA Serbinova and L Packer. Antioxidant properties of alpha-tocopherol and alpha-tocotrienol. *Methods Enzymol.* 1994; **234**, 354–366
- [55] G Horvath, L Wessjohann, J Bigirimana, M Jansen, Y Guisez, R Caubergs, and N Horemans. Differential distribution of tocopherols and tocotrienols in photosynthetic and non-photosynthetic tissues. *Phytochemistry.* 2006, **67**, 1185–1195
- [56] NN Okoye, DL Ajaghaku, HN Okeke, EE Ilodigwe, CS Nworu, and FB Okoye. β -Amyrin and α -amyrin acetate isolated from the stem bark of *Alstonia boonei* display profound anti-inflammatory activity. *Pharm. Biol.* 2014; **52(11)**, 1478-1486.
- [57] VD Ebajo Jr., CC Shen, Y Consolacion and CY Ragasa. Terpenoids and sterols from *Hoya multiflora* Blume. *J. Appl. Pharm. Sci.* 2015; **5(4)**, 33-039.
- [58] MLL De Amorim, WM Godinho, FC Archanjo and CFF Graell. Chemical constituents of *Pseudobrickellia brasiliensis* leaves (Spreng.) R.M. King and H. Rob. (*Asteraceae*). *Rev. Bras. Plant Med.* 2016; **18(2)**, 408-414.
- [59] MF King, LC Boyd and BW Sheldon. Antioxidant properties of individual phospholipids in a salmon oil model system. *J. Am. Oil Chem. Soc.* 1992; **69**,545-551.
- [60] T Koga, and J Terao. Phospholipids increase radical-scavenging activity of vitamin E in a bulk oil model system. *J. Agric. Food Chem.* 1995; **43**, 1450-1454
- [61] FJ Hidalgo, MM Leon and R Zamora. Antioxidative activity of amino phospholipids and phospholipid/amino acid mixtures in edible oils as determined by the Rancimat method. *J. Agric. Food Chem.* 2006; **54**, 5461-5467.
- [62] RD Reichert. Oilseed medicinal: in natural drugs, dietary supplements and in new functional foods. *Trends Food Sci. Technol.* 2002; **13**, 353–60.

- [63] KM Al-Ismail, AK. Alsaed, R Ahmad and M Al-Dabbas. Detection of olive oil adulteration with some plant oils by GLC analysis of sterols using polar column. *Food Chem.* 2010; **121**, 1255–1259.
- [64] NG Kammoun, and W Zarrouk. Exploratory chemometric analysis for the characterisation of Tunisian olive cultivars according to their lipid and sterolic profiles. *Int. J. Food Sci. Technol.* 2012; DOI:10.1111/j.1365-2621.2012.02997.x
- [65] WM Al-Rousan, RY Ajo, KM. Al-Ismail, A Attlee, RR Shaker and TM Osaili. Characterization of acorn fruit oils extracted from selected Mediterranean *Quercus* species. *Grasas Aceites* 2013; **64(5)**, 554-560.
- [66] H Schaller. The role of sterols in plant growth and development. *Prog. Lipid Res.* 2003; **42**, 163–175.
- [67] MA Hartmann. Plant sterols and the membrane environment. *Trends Plant Sci.* 1998, **3(5)**, 170-175.
- [68] REJ Ostlund, SB Racette, and WF Stenson. Effects of trace components of dietary fat on cholesterol metabolism: phytosterols, oxysterols, and squalene. *Nutr. Rev.* 2002, 60:349–59.
- [69] M Law. Plant sterol and stanol margarines and health. *Br. Med. J.* 2000; **320**,861–4.
- [71] OS Ijarotimi and OO Keshinro. Comparison between the amino acid, fatty acid, mineral and nutritional quality of raw, germinated and fermented African locust bean (*Parkia biglobosa*) flour. *Acta Sci. Pol. Technol. Alim.* 2012; **11(2)**, 151-165
- [72] Salunkhe DK, Kadam SS (1989) Handbook of world food legumes, nutritional chemistry, processing technology and utilization. Boca Raton, FL: CR Press.
- [73] MA Mune-Mune, SR Minka, IL Mbome and F-X Etoa. Nutritional potential of Bambara bean protein concentrate. *Pak J. Nutr.* 2011; **10(2)**, 112-119.
- [74] Food and Agriculture Organization of the United Nations (FAO), (2013). Dietary protein quality evaluation in human nutrition, FAO food and nutrition paper No. 92, Food and Agriculture Organization of the United Nations Viale delle Terme di Caracalla, Rome.
- [75] ENV/JM/MONO (2001)15. Unclassified. 2001. Series on the Safety of Novel Foods and Feeds No.2, Consensus Document on Compositional Considerations for New Varieties of Soybean: Key Food and Feed Nutrients and Anti-nutrients, 30-November-2001